

Contents

Preface

Notation

- 1. Introduction** **Page 1**
 - 1.1 Expectations
 - 1.2 The problem to be solved
 - 1.3 Data sets
 - 1.4 Software

- 2. Why a statistical approach?** **Page 7**
 - 2.1 Investigating the sample data
 - 2.2 Measures of central tendency
 - 2.3 Measures of spread or variability
 - 2.4 Graphical descriptions of the data
 - 2.5 Other useful descriptive statistics
 - 2.6 Discrete data
 - 2.7 Into the unknown
 - 2.8 Worked examples
 - 2.8.1 Coal project data, calorific values
 - 2.8.2 Iron ore example
 - 2.8.3 Wolfcamp data
 - 2.8.4 Scallops, total catch
 - 2.9 Exercises

- 3. Normal (Gaussian) distributions** **Page 29**
 - 3.1 The gap between data and population
 - 3.2 Is it a Normal distribution?
 - 3.3 Estimating population parameters
 - 3.3.1 estimating the population average

3.3.2	estimating the standard deviation	
3.3.3	confidence intervals for standard deviation	
3.3.4	confidence intervals for mean	
3.4	Selection (grade/tonnage) calculations	
3.5	Summary of chapter	
3.6	Worked examples	
3.6.1	Coal project, calorific values	
3.6.2	Iron ore example	
3.6.3	Wolfcamp data	
3.6.4	Scallop data, total catch	
3.7	Exercises	
4.	Lognormal distributions (and others)	Page 63
4.1	The lognormal distribution	
4.1.1	estimating the mean of a lognormal population	
4.1.2	confidence intervals on the population mean	
4.2	The three parameter lognormal	
4.3	Selection (grade/tonnage) calculations	
4.3.1	two parameter lognormal — reef widths	
4.3.2	three parameter lognormal — gold grades	
4.4	More complex distributions and mixtures	
4.4.1	mixtures of Normal or lognormal populations	
4.5	Worked examples	
4.5.1	Scallops, total caught	
4.5.2	Organic matter in soil	
4.5.3	Calcium in limestone	
4.5.4	Geevor Tin mine, Cornwall	
4.6	Exercises	
5.	Discrete distributions	Page 99
5.1	Review of Discrete Moments	
5.2	Bernoulli and Binomial Distributions	
5.3	Negative Binomial and Geometric Distributions	
5.4	Poisson Distribution	
5.5	Mixtures of Poisson Distributions (Compound Poisson)	
5.5.1	Oswego Zircon data	
5.5.2	Other examples	

5.6 Spatial Considerations

5.7 Solved Problems

5.8 Exercises

6. Hypothesis testing **Page 129**

6.1 Single sample tests

6.1.1 test on sample mean

6.1.2 test on sample standard deviation

6.2 Two sample tests

6.2.1 test on standard deviations

6.2.2 test on means

6.2.3 paired sampling

6.2.4 test for sample distribution

6.3 Worked examples

6.3.1 Heights of students

6.3.2 Geevor tin mine – development versus stope

6.4 Exercises

7. Relationships **Page 141**

7.1 Straight line relationships

7.1.1 quantifying the strength of the relationship

7.1.2 Predicting one variable from the other

7.1.3 Calorific Value versus Ash Content

7.1.4 Calorific Value versus Sulphur Content

7.2 Other worked examples

7.2.1 Gold grade versus reef width

7.2.2 Scallops caught

7.2.3 Application — Krige's Regression Effect

7.3 Relationships involving more than two variables

7.3.1 Predicting Sulphur from Calorific Value and Ash Content

7.3.2 Application — Krige's moving average template

7.3.3 Curvilinear Regression

7.3.4 Application — Polynomial Trend Surface Analysis

7.4 Exercises

8. The spatial element **Page 177**

8.1 Including location as well as value

- 8.2** Spatial relationships
- 8.3** Inverse distance estimation
- 8.4** Worked examples
 - 8.4.1 Coal project, calorific values
 - 8.4.2 Iron ore project
 - 8.4.3 Wolfcamp data
 - 8.4.4 Scallops caught
- 8.5** Exercises

9. The semi-variogram

Page 197

- 9.1** The experimental semi-variogram
 - 9.1.1 Irregular sampling
 - 9.1.2 Cautionary notes
- 9.2** Modelling of the semi-variogram function
 - 9.2.1 The linear model
 - 9.2.2 The generalised linear model
 - 9.2.3 The Spherical model
 - 9.2.4 The exponential model
 - 9.2.5 Gaussian model
 - 9.2.6 The hole effect model
 - 9.2.7 Paddington mix model
 - 9.2.8 Judging how well the model fits the data
 - 9.2.9 equivalence to covariance function
 - 9.2.10 the nugget effect
- 9.3** Worked examples
 - 9.3.1 Silver example from Practical Geostatistics 1979
 - 9.3.2 Coal project: calorific values
 - 9.3.3 Wolfcamp aquifer
- 9.4** Exercises

10.1 Estimation error

- 10.1.1 one sample estimation
- 10.1.2 another single sample
- 10.1.3 two sample estimation
- 10.1.4 another two sample estimation
- 10.1.5 three sample estimator

10.2 Choosing the optimal weights

- 10.2.1 three sample estimation
- 10.2.2 the general form for the ‘optimal’ estimator
- 10.2.3 confidence levels and degrees of freedom
- 10.2.4 simple kriging

10.3 Ordinary kriging

- 10.3.1 ‘optimal’ unbiased estimator
- 10.3.2 alternate form: matrices
- 10.3.3 alternate form: covariance
- 10.3.4 three sample estimation

10.4 Worked examples

- 10.4.1 Coal project, calorific values
- 10.4.2 Iron ore example, (Page95)
- 10.4.3 Wolfcamp, residuals from quadratic surface

10.5 Cross validation

- 10.5.1 cross cross validation

10.6 Exercises

11.1 The impact on the distribution

- 11.1.1 Iron ore example, Normal example
- 11.1.2 Geevor Tin Mine, lognormal(ish) example

11.2 The impact on kriging

- 11.2.1 the use of auxiliary functions
- 11.2.2 Iron ore example, Page 95
- 11.2.3 Wolfcamp aquifer, quadratic residuals

12. Other kriging approaches	Page 301
12.1 Universal kriging	
12.1.1 Wolfcamp aquifer	
12.2 Lognormal kriging	
12.2.1 the proportional effect	
12.2.2 the lognormal transformation	
12.2.3 Geevor Tin Mine, grades	
12.2.4 SA Gold Mine	
12.3 Indicator kriging	
12.4 Rank uniform kriging	
12.5 Summary of chapter	
13. Bibliography	Page 325
Tables	Page 329
Data Sets	Page 373
Index	Page 409